

#9/BM
13-26-97

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

Vuorinen et al

Atty. Ref.: 30-336

Serial No. 08/542,646

Group: 1303

Filed: October 13, 1995

Examiner: Alvo

For: Method of Treating Cellulose Pulp

Honorable Commissioner of
Patents and Trademark
Washington, DC 20231

EVIDENTIARY DECLARATION OF TAPANI VUORINEN

Sir:

I, Tapani Vuorinen, hereby say and declare:

1. My curriculum vitae is attached hereto as Exhibit A. I am one of the co-inventors of the above-identified application.

2. The Lachenal et al reference (1982 International Bleaching Conference, pages 145-151) of record in this application describes a combination of an acidic treatment with a subsequent peroxide bleaching stage. The charge of sulfuric acid in the acidic stage is 3% on pulp which, according to the authors, corresponded to a pH-2. In reality the pH must have been lower. The pH can be evaluated from the charge of the acid when the anionic charge of the pulp is approximately known (Laine et al, Colloids and Surfaces, A, 88, 1994, 277-287). Such a calculation shows that the pH corresponding to a 3% charge of sulfuric acid is -1.5. At the 12% pulp consistency pH -2.0 would be obtained with a 1.3% charge of sulfuric acid.

VUORINEN et al
Serial N . 08/542,646

- 2 -

3. Lachenal et al show that a major effect of their acidic treatment, irrespective of its temperature, is a reduction in manganese content of the pulp. Because manganese catalyzes decomposition of peroxide, a better bleaching result was obtained with the acidic pretreatment. It may be noted that peroxide bleaching is currently widely applied for bleaching of kraft pulps but an acidic pretreatment is not generally used for the removal of manganese ions. This is mainly because efficient manganese removal requires a high acid charge and a low pH (≤ 2). Instead of a treatment with acid, pulps are usually treated with complexing agents in order to remove manganese or other heavy metal ions before a peroxide bleaching stage.

4. Lachenal et al also show that the efficiency of the acid treatment is increased when it is performed at elevated temperatures. Experiments were done at 50, 70, and 90°C. In the conclusions the temperature range was restricted to 60-80°C. No specific reason was given for why a treatment at 90°C was not in the preferred range, however my understanding is that at 90°C the high acid charge, corresponding to a pH of ~ 1.5 , decreases the viscosity of the pulp dramatically. The strength properties become poor which makes the treatment under these conditions impractical. In fact, Lachenal et al describe viscosities and strength properties only for pulps treated with the acid at 70°C where the hydrolysis of cellulose must have been much less extensive.

5. The above-identified application relates to a method for selectively hydrolysing hexenuronic acid groups in pulps. Before the invention the existence and importance of hexenuronic acid as related to bleaching chemicals was not, to my knowledge, known in the art. Hexenuronic acid groups consume bleaching

VUORINEN et al
Serial N . 08/542,846

- 3 -

chemicals, bind heavy metal ions, cause brightness reversion, and form a major oxalate source in pulp bleaching. For these reasons it is often advantageous to remove the hexenuronic acid groups before final bleaching.

6. The hydrolysis of hexenuronic acid groups follows an exceptional reaction kinetics. The reaction rate gets its maximum value at pH 3-4 (Vuorinen et al, 1996 International Pulp Bleaching Conference, 48-51). The reaction cannot be speeded up much by lowering the pH further to 1.5-2. In order to remove most of the hexenuronic acid groups a relatively high temperature and long treatment time is necessary. Typically the temperature is over 85°C, and preferably at least 90°C, and the treatment time several hours.

7. In 1995 I, with the cooperation of others, carried out extensive studies on how the selective hydrolysis of hexenuronic acid groups could be implemented in bleaching of kraft pulps. We were fully aware of the negative effects of too low a pH and therefore we were also looking for an optimum pH for treatment. The viscosity decreased least at pH 3-4, preferably 3-3.5 (depending on the pulp). Since these studies I have continuously been involved in development and application of the selective hydrolysis stage. My experience is that the viscosity of pulp is really a factor that must be under control although the hydrolysis is carried out at pH 3-3.5.

8. A significant difference between the acidic treatments of Lachenal et al and the invention is illustrated with the following examples. In Table 1 (appended as Exhibit B) softwood kraft pulp (kappa number 25.4) was treated at pH 2 (0.01 M sulfuric acid) and 3.5 (formate buffer) at 90°C. Both treatments reduced the kappa number by 3.6-3.7 units (Table 1). Simultaneously 32-38 meq/kg of hexenuronic

VUORINEN et al**- 4 -****Serial No. 08/542,846**

acid groups were removed. This reduction corresponds to a 3.4 units decrease in the kappa number (Vuorinen et al, 1996 International Pulp Bleaching Conference, 43-51). Thus, the kappa number reductions in the acidic stages were solely caused by reductions in the hexenuronic acid content of the pulp. As Table 1 shows, at pH 3.5 the viscosity was decreased but at pH 2 it dropped dramatically. This example shows that the treatment described by Lachenal et al at 90°C is impractical. A key is that the pH of the acidic stage must be high enough in order to prevent the hydrolysis of cellulose.

9. In a second example (Table 2, Exhibit C hereto) an oxygen bleached hardwood kraft pulp was treated at pH 1.5 (2% sulfuric acid on pulp) and 3.5 (formiate buffer) at 90°C. The kappa number reduction was in both cases 4.2-4.5 units (Table 2). Because the decrease in the hexenuronic acid content was 29-31 meq/kg, a small fraction of the kappa number reduction could be caused by hydrolysis of lignin structure. The viscosity losses were moderate at pH 3.5 but at pH 1.5 the viscosity again dropped dramatically. Thus, the conclusions are the same as from the previous example.

10. In a third example (Table 3, Exhibit D), unbleached and oxygen bleached hardwood kraft pulps were treated at pH 2 (1.3% sulfuric acid on pulp) and pH 3.5 (formiate buffer or adjustment with sulfuric acid) at 70 and 90°C. After the treatments the pulps were neutralized and treated with a complexing agent (0.2% EDTA on pulp) in order to remove heavy metal ions. Finally the pulps were bleached with alkaline hydrogen peroxide (3% on pulp) for two hours at 90°C. As Table 3 shows, removal of hexenuronic acid groups at 70°C was limited to 2.4-6.5

VUORINEN et al
Serial No. 08/542,646

- 5 -

meq/kg. The kappa number reduction under the same conditions was 0.7-1.1 units. At 90°C the removal of hexenuronic acid groups was much more extensive which also resulted in a lower kappa number after the acidic stage. None of the acidic pretreatments improved lignin removal in the peroxide stage. This can clearly be seen by comparing the kappa number difference before and after the peroxide stage. Thus, at least in this case, the effect of the acidic pretreatment was restricted to the removal of hexenuronic acid groups.

11. A last example (Table 4, Exhibit E) shows that the effects described by Lachenal et al for an acidic pretreatment at pH ~ 2 at 60-80°C lack generality. To the contrary, the selective hydrolysis described according to the invention gives effects that can be predicted from the hexenuronic acid content of the pulp.

12. In conclusion, the acidic treatments described by Lachenal et al and the invention are clearly different. In the first case the treatment is optimally at pH ~2 at 60-80°C. The low pH prevents practical treatments at higher temperatures where the removal of hexenuronic acid groups becomes prominent. The bleaching result depends on the pulp. In some cases the acidic pretreatment gives no advantage.

13. The acidic treatment of the invention is optimally performed at about pH 3-4 and above 85°C, preferably at or above 90°C. The effect of the treatment is based on the removal of hexenuronic acid groups and is thus predictable from their content in the pulp.

14. I declare further that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be

VUORINEN et al**- 6 -****Serial No. 08/548,646**

true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Tapani Vuorinen

Tapani Vuorinen

Date: March 6, 1997

Jouko Tapani Vuorinen

Curriculum vitae

CURRICULUM VITAE**Personal data**

Full name: Vuorinen, Jouko Tapani
Date and place of birth: 22 September 1957, Turku
Parents: Lauri Vuorinen and Aila Vuorinen, b. Kyrö
Married: 1985 with Kristina Virtanen
Children: Pauliina Mikaela, 1990, Heidi Adellina, 1995

Education

Matriculation Examination, Unsikaupunki High School, 29 May 1976
Master of Science in Technology, Helsinki University of Technology, 29 August 1980
Doctor of Technology, Helsinki University of Technology, 30 August 1988

Professional appointments

Assistant in Wood Chemistry, Helsinki University of Technology, 1.1.1980-31.7.1991
Senior Research Fellow, Academy of Finland, 1.8.1991-31.3.1996.
Docent in Organic Chemistry, Turku University, 2.3.1994-
Docent in Forest Products Chemistry, Helsinki University of Technology, 1.6.1994-
Associate Professor in Forest Products Chemistry, Helsinki University of Technology,
1.4.1996-

International activities

University of Notre Dame, USA, visiting scientist, 22.8.1988-28.2.1989, 18.9.-19.11.1989.
International Carbohydrate Organization, member, 1994-

Other activities

Secretary, National program on carbohydrates in process technology (Technology
Development Centre Finland), 1993-
Lecturer, Turku University
 Synthetic organic and bio-organic chemistry: synthetic carbohydrate chemistry, 1993
 Wood Chemistry, 1994
Lecturer, Helsinki University of Technology
 Methods in chemistry of forest products technology, 1993-
 Chemistry of lignin and its properties in pulping processes, 1993-

Memberships in scientific societies

Finnish Chemical Society, 1980-
American Chemical Society, Division of Carbohydrate Chemistry, Division of Agricultural
& Food Chemistry, Division of Cellulose, Paper & Textile Chemistry, 1989-

Awards and honours

Honorary award of young scientists, Finnish Chemical Society, 1980

Special mention, Technology Development Centre Finland, National program on carbohydrates in process technology, 1993

Honorary mention, National Blue Globe contest for inventions promoting clean environment, 1996

List of publicationsPapers in scientific journals

1. Tapani Vuorinen and Eero Sjöström, Kinetics of alkali-catalyzed isomerization of D-glucose and D-fructose in ethanol-water solutions, *Carbohydrate Research*, 108 (1982) 23-29.
2. Tapani Vuorinen, Kinetics of alkaline degradation of maltose in ethanol-water solutions, *Carbohydrate Research*, 108 (1982) 213-219.
3. Tapani Vuorinen and Eero Sjöström, The effect of solvent on the oxygen oxidation of polysaccharides, *Journal of Wood Chemistry and Technology*, 2 (1982) 129-145.
4. Tapani Vuorinen, Alkali-catalyzed oxidation of D-glucose with sodium 2-anthraquinonesulfonate in ethanol-water solutions, *Carbohydrate Research*, 116 (1983) 61-69.
5. Tapani Hyppänen, Eero Sjöström and Tapani Vuorinen, Gas-liquid chromatographic determination of hydroxy carboxylic acids on a fused-silica capillary column, *Journal of Chromatography*, 261 (1983) 320-323.
6. Tapani Vuorinen, Cleavage of D-arabino-hexos-2-ulose and glyoxal with hydrogen peroxide, *Carbohydrate Research*, 127 (1984) 319-325.
7. Tapani Vuorinen, Alkali-catalyzed rearrangement of glyoxal and 3-deoxy-D-erythro-hexos-2-ulose, *Carbohydrate Research*, 127 (1984) 327-330.
8. Tapani Vuorinen, Effect of calcium ion on rearrangement of aldoses-2-uloses to aldonic acids, *Carbohydrate Research*, 133 (1984) 329-333.
9. Tapani Vuorinen, 2-C-Carboxyaldoses and aldonic acids from cellobiose, maltose, and 4-O-methyl-D-glucose with 2-anthraquinonesulfonic acid, *Carbohydrate Research*, 141 (1985) 307-317.

10. Tapani Vuorinen, Cleavage of the intermediate hydroperoxides in the oxidation of D-glucose and D-fructose with oxygen, *Carbohydrate Research*, 141 (1985) 319-322.
11. Tapani Vuorinen, Mechanisms and kinetics of isomerization, degradation, and oxidation of reducing carbohydrates; reaction paths in alkaline solutions containing oxygen and 2-anthraquinonesulfonic acid, Dissertation, *Annales Academiae Scientiarum Fennicae, Series A, II. Chemica*, 220 (1988) 160 pp.
12. Tapani Vuorinen, Stabilizing interactions between a free carbonyl group and hydroxyl groups in carbohydrates, *Annales Academiae Scientiarum Fennicae, Series A, II. Chemica*, 227 (1990) 191-194.
13. Jian Wu, Anthony S. Serianni, and Tapani Vuorinen, Furanose ring anomerization: kinetic and thermodynamic studies of the D-2-pentuloses by ^{13}C -n.m.r. spectroscopy, *Carbohydrate Research*, 206 (1990) 1-12.
14. Tapani Vuorinen and Anthony S. Serianni, ^{13}C -Substituted pentos-2-uloses: synthesis and analysis by ^1H - and ^{13}C -n.m.r. spectroscopy, *Carbohydrate Research*, 207 (1990) 185-210.
15. Tapani Vuorinen and Anthony S. Serianni, Synthesis of D-erythro-2-pentulose and D-threo-2-pentulose and analysis of the ^{13}C - and ^1H -n.m.r. spectra of the 1- ^{13}C - and 2- ^{13}C -substituted sugars, *Carbohydrate Research*, 209 (1990) 13-31.
16. Anthony S. Serianni, Tapani Vuorinen, and Paul B. Bondo, Stable isotopically-enriched D-glucose: strategies to introduce carbon, hydrogen and oxygen isotopes at various sites, *Journal of Carbohydrate Chemistry*, 9 (1990) 513-541.
17. Tapani Vuorinen, Tapani Hyppänen, and Eero Sjöström, Oxidation of D-glucose with oxygen in alkaline methanol-water mixtures: a convenient method of producing crystalline sodium D-arabinonate, *Starch/Stärke*, 43 (1991) 194-198.
18. Jian Wu, Paul B. Bondo, Tapani Vuorinen, and Anthony S. Serianni, ^{13}C - ^{13}C Spin coupling constants in aldoses enriched with ^{13}C at the terminal hydroxymethyl carbon: effect of coupling pathway structure on J_{CH} in carbohydrates, *Journal of the American Chemical Society*, 114 (1992) 3499-3506.
19. Tapani Vuorinen, The role of carbohydrates in anthraquinone alkali pulping, *Journal of Wood Chemistry and Technology*, 13 (1993) 97-125.
20. Anita Teleman, Vesa Harjunpää, Maija Tenkanen, Johanna Buchert, Tiina Hausalo, Torbjörn Drakenberg and Tapani Vuorinen, Characterization of 4-deoxy- β -L-threo-hex-4-enopyranosyluronic acid attached to xylan in pine kraft pulp and pulping liquor by ^1H and ^{13}C NMR spectroscopy, *Carbohydrate Research*, 272 (1995) 55-71.

21. Anita Teleman, Tiina Hausalo, Maija Tenkanen and Tapani Vuorinen, Identification of the acidic degradation products of hexenuronic acid by NMR spectroscopy, *Carbohydrate Research*, 280 (1996) 197-208.
22. Johanna Buchert, Anita Teleman, Vesa Harjunpää, Maija Tenkanen, Liisa Viikari and Tapani Vuorinen, Effect of cooking and bleaching on the structure of xylan in conventional pine kraft pulp, *Tappi Journal*, 78 (1995):1, 125-130.
23. Kenneth N. Drew, Timothy J. Church, Bisida Basu, Tapani Vuorinen and Anthony S. Serianni, L-(1-¹³C)- and (2-¹³C)ascorbic acid: synthesis and NMR characterization, *Carbohydrate Research*, 284 (1996) 135-143.
24. Anna Suurnäkki, Tie-Qiang Li, Johanna Buchert, Maija Tenkanen, Liisa Viikari, Tapani Vuorinen and Lars Ödberg, Effects of enzymatic removal of xylan and glucomannan on the pore size distribution of kraft fibres, *Holzforschung*, 51 (1997) 27-33.

Papers at scientific meetings

25. Tapani Vuorinen and Eero Sjöström, Selective oxidation and stabilization of carbohydrates with oxygen, *The Ekman-Days 1981, Organic & Physical Chemistry of Pulp and Bleaching Processes - New Development*, Vol. 2, International Symposium of Wood and Pulp Chemistry, Stockholm, June 9-12, 1981, 90-93.
26. Tapani Vuorinen, Oxidation of reducing carbohydrates by oxygen, 14th International Carbohydrate Symposium, Stockholm, August 14-19, 1988, Abstract No. D 15.
27. Tapani Vuorinen and Anthony S. Serianni, Synthesis of (¹³C)-enriched pentos-2-uloses and their analysis by ¹H and ¹³C NMR spectroscopy, 198th ACS National Meeting, Miami Beach, September 10-15, 1989, Division of Carbohydrate Chemistry, Abstract No. 53.
28. Tapani Vuorinen and Anthony S. Serianni, Synthesis of (¹³C)-enriched 2-pentuloses and their characterization by ¹H (620 MHz) and ¹³C (75 MHz) NMR spectroscopy, 198th ACS National Meeting, Miami Beach, September 10-15, 1989, Division of Carbohydrate Chemistry, Abstract No. 54.
29. Tapani Vuorinen, Degradation of aldoses, ketoses, and their derivatives with hydrogen peroxide and performic acid, 201st ACS National Meeting, Atlanta, April 14-19, 1991, Division of Carbohydrate Chemistry, Abstract No. 63.
30. Tapani Vuorinen, Synthesis of xylitol from D-glucose; a literature survey and preliminary studies, 6th European Symposium on Carbohydrate Chemistry, Edinburgh, September 8-13, 1991, Abstract No. B 89.
31. Tapani Vuorinen, Monika Blättler, Tarja Hulkkonen, Minna Pyykkönen and Mari Takanen, D-Arabinitol by hydrogenation of D-arabinonic acid, 2nd International

- Workshop Carbohydrates as Organic Raw Materials, Lyon, July 2-3, 1992, Abstract No. C9.
32. Jianju Luo and Tapani Vuorinen, Stabilities of 1,4- and 1,5-lactones and structure-reactivity relationships in their hydrolysis; a comprehensive study with all hexono-, pentono- and tetronolactones, 16th International Carbohydrate Symposium, Paris, July 5-10, 1992, Abstract No. C18.
 33. Tapani Vuorinen, Aldos-2-uloses; structure and reactions in solution, 7th European Symposium on Carbohydrate Chemistry, Cracow, August 22-27, 1993, Abstract No. B 3.
 34. Tapani Vuorinen, Hiilihydraattikemian tutkimuksesta avain uuteen keitto- ja valkaisu-tekniikkaan (Research on carbohydrate chemistry as a key to new pulping and bleaching technology), Tulevaisuuden sellutekniikka-seminaari, Helsinki, January 11, 1994.
 35. Nina Engström, Anne Vikkula, Anita Teleman and Tapani Vuorinen, Puun hemiselluloosien fysikaalinen käyttäytyminen delignifioinnin yhteydessä - konventionaalinen sulfaattikeitto (Physical behaviour of wood hemicelluloses during delignification - conventional kraft pulping), Tulevaisuuden sellutekniikka-seminaari, Helsinki, January 11, 1994.
 36. Anita Teleman, Vesa Harjunpää, Maija Tenkanen and Tapani Vuorinen, Characterization by ^1H and ^{13}C n.m.r. spectroscopy of hexenuronic acid, degradation product of 4-O-methylglucuronic acid, found in pulp and pulp liquor, Tulevaisuuden sellutekniikka-seminaari, Helsinki, January 11, 1994.
 37. Johanna Buchert, Maija Tenkanen, Anita Teleman, Vesa Harjunpää, Jari Tuominen, Tapani Vuorinen, Janne Laine, Raimo Alén, Tiina Hausalo and Tarja Tamminen, Sellumassojen karboksyyliyhdyntien karakterisointi (Characterization of carboxyl groups in cellulose pulps), Tulevaisuuden sellutekniikka-seminaari, Helsinki, January 11, 1994.
 38. Anita Teleman, Vesa Harjunpää, Torbjörn Drakenberg, Maija Tenkanen and Tapani Vuorinen, Characterization by ^1H and ^{13}C NMR spectroscopy of hexenuronic acid found in pulp and pulp liquor, 12th European Experimental NMR Conference, Oulu, June 5-10, 1994, 233.
 39. Nina Engström, Anita Teleman and Tapani Vuorinen, Changes in chemical structure of wood xylans during kraft pulping, Plant Polysaccharide Symposium, Guelph, July 13-15, 1994, P52.
 40. Anita Teleman, Vesa Harjunpää, Torbjörn Drakenberg, Maija Tenkanen and Tapani Vuorinen, Characterization by ^1H and ^{13}C NMR spectroscopy of hexenuronic acid linked to xylan in kraft pulps, 17th International Carbohydrate Symposium, Ottawa, July 17-22, 1994, 81.

41. Nina Engström, Anita Teleman and Tapani Vuorinen. Dissolved hemicelluloses in kraft pulping of pine wood, 17th International Carbohydrate Symposium, Ottawa, July 17-22, 1994, 95.
42. Johanna Buchert, Anita Teleman, Vesa Harjunpää, Maija Tenkanen, Liisa Viikari and Tapani Vuorinen, Effect of cooking and bleaching on the surface chemistry of conventional pine kraft pulp, 3rd European Workshop on Lignocellulosics and Pulp, Stockholm, August 29-31, 1994, 16-19.
43. Tapani Vuorinen, Hydrogenation of aldonolactones - reaction mechanism, 3rd International Workshop on Carbohydrates as Organic Raw Materials, Wageningen, November 28-29, 1994.
44. Anita Teleman, Vesa Harjunpää, Maija Tenkanen, Johanna Buchert, Torbjörn Drakenberg and Tapani Vuorinen, Characterization of hexenuronic acid attached to xylan in pine kraft pulp and pulping liquor by ^1H and ^{13}C NMR spectroscopy, International Meeting on Carbohydrate Structure, London, December 19-20, 1994, 2.
45. Tapani Vuorinen, Modernin hiilihydraattianalytiikan tarjoamat mahdollisuudet puunjälöstukseen liittyvässä tutkimuksessa (Possibilities of modern carbohydrate analysis in forest products research), Analyytinen seminaari, KCL, Espoo, February 8, 1995, 32-46.
46. Tapani Vuorinen, Possibilities and challenges for modern carbohydrate research in wood and pulping chemistry, 8th International Symposium on Wood and Pulping Chemistry, Helsinki, June 6-9, 1995, Proceedings, Vol. I, 223-230.
47. Johanna Buchert, Anita Teleman, Gilbert Carlsson, Maija Tenkanen, Janne Laine, Vesa Harjunpää, Tiina Hausalo, Liisa Viikari, Göran Ström and Tapani Vuorinen, Effect of pulping and bleaching on the surface composition of kraft pulps, 8th International Symposium on Wood and Pulping Chemistry, Helsinki, June 6-9, 1995, Proceedings, Vol. I, 567-573.
48. Nina Engström, Anne Vikkula, Anita Teleman and Tapani Vuorinen, Structure of hemicelluloses in pine kraft cooking liquors, 8th International Symposium on Wood and Pulping Chemistry, Helsinki, June 6-9, 1995, Proceedings, Vol. III, 195-200.
49. Anita Teleman, Tiina Hausalo, Maija Tenkanen and Tapani Vuorinen, Identification of the acidic degradation products of hexenuronic acid by NMR spectroscopy, 8th International Symposium on Wood and Pulping Chemistry, Helsinki, June 6-9, 1995, Vol. III, 109-114.
50. Tarja Tamminen, Tapani Vuorinen, Maija Tenkanen, Tiina Hausalo and Bo Hertilg, Analysis of lignin and lignin-carbohydrate complexes isolated from black liquor, 8th International Symposium on Wood and Pulping Chemistry, Helsinki, June 6-9, 1995, Proceedings, Vol. II, 297-302.

-
51. Mia Mitikka, Raivo Teeäär, Maija Tenkanen, Janne Laine and Tapani Vuorinen, Sorption of xylans on cellulose fibers, 8th International Symposium on Wood and Pulping Chemistry, Helsinki, June 6-9, 1995, Proceedings, Vol. III, 231-236.
 52. Johanna Buchert, Maija Tenkanen, Monika Ek, Anita Teleman, Liisa Viikari and Tapani Vuorinen, Effects of pulping and bleaching on pulp carbohydrates and technical properties, International Pulp Bleaching Conference, Washington D.C., April 14-18, 1996, TAPPI Proceedings, 39-42.
 53. Tapani Vuorinen, Johanna Buchert, Anita Teleman, Maija Tenkanen and Pirjo Fagerström, Selective hydrolysis of hexenuronic acid groups and its application in ECF and TCF bleaching of kraft pulps, International Pulp Bleaching Conference, Washington D.C., April 14-18, 1996, TAPPI Proceedings, 43-51.
 54. Tapani Vuorinen, Heksenuronihapporyhmien selektiiviseen poistoon perustuva uusi valkaisuvaihe (a new bleaching stage based on selective removal of hexenuronic acid groups), Suomen Paperi-insinöörien Yhdistyksen vuosikokous, Helsinki, 24-25.4.1996.
 55. Erkki Räsänen and Tapani Vuorinen, Removal of hexenuronic acid groups from kraft pulps by selective hydrolysis, Plant Polysaccharide Symposium, Nantes, July 17-19, 1996, 145.
 56. Tapani Vuorinen, Bo Hortling, Johanna Buchert, Maija Tenkanen and Liisa Viikari, National programme on wood hemicelluloses finds applications in pulp and paper technology, XVIII International Carbohydrate Symposium, Milano, July 21-26, 1996, 628.
 57. Tapani Vuorinen, Uusi esivalkaisuvaihe heksenuronihapporyhmien selektiiviseen poistoon (a new prebleaching stage for selective removal of hexenuronic acid groups), AEL-METSKO, Insko-seminaarit, Pori, 26-27.11.1996, s. XI 1-20.
 58. Tapani Vuorinen, Behaviour of wood hemicelluloses during pulping and bleaching, Workshop on chemistry influencing fiber structure and properties, Concerted Action in Wood Chemistry, February 6-7, 1997.

Scientific reports

59. Tapani Vuorinen, Ksylitolin tuottaminen D-glukoosista (Production of xylitol from D-glucose), report to the Academy of Finland, February 27, 1991.
60. Tapani Vuorinen, Production of xylitol from D-glucose, Intermediate report, October 8, 1992.

-
61. Tapani Vuorinen, Hydrogenation of D-arabinono-1,4-lactone, Intermediate report, September 9, 1993.
 62. Tapani Vuorinen, Isomerization of D-arabinitol, Intermediate report, September 27, 1993.
 63. Tapani Vuorinen (toim.), Hiilihydraatit prosessiteollisuudessa (Carbohydrates in process technology), Teknologiaohjelmaraaportti 9/96, Teknologian kehittämiskeskus, Helsinki, 1996, 48 p.

Patents

64. Eero Sjöström ja Tapani Vuorinen, Menetelmä hiilihydraattien selektiiviseksi hapettamiseksi ja stabiiloimiseksi happipitoisella kaasulla (Method for selective oxidation and stabilization of carbohydrates with oxygen containing gas), FI 62356, 31.8.1992.
65. Amylum N.V., Tapani Vuorinen (inventor), Production of xylitol, PCT/BE93/00013.
66. Matti Siika-aho, Johanna Buchert, Tapani Vuorinen, Anita Teleman, Maija Tenkanen, Michael Bailey and Liisa Viikari, Method and enzymatic preparation for treatment of cellulose pulps, PCT/FI95/00317, 5.6.1995.
67. Tapani Vuorinen, Johanna Buchert, Anita Teleman and Maija Tenkanen (keks.) A. Ahlström co., Menetelmä selluloosamassojen käsittelemiseksi (Method for treatment of cellulose pulps), FI 944808, 13.10.1994. PCT application and national applications in several countries.

March 3, 1997

Tapani Vuorinen

EXHIBIT B

Table 1. Effect of acidic treatments on the properties of a softwood kraft pulp. Δ HexA denotes the amount of hexamuronic acid groups removed in the acid treatment. Treatments were carried out at 2 % pulp consistency.

pH	Time (h)	Kappa number	Viscosity (ml/g)	Δ HexA (msq/kg)
-	-	25.4	1166	-
2	1	22.5	1074	18.7
2	2	22.1	962	25.7
2	4	21.7	869	32.8
3.5	1	23.3	1107	13.6
3.5	2	22.8	1117	20.4
3.5	4	21.8	1069	32.2

EXHIBIT C

Table 2. Effect of acidic treatments on the properties of an oxygen bleached hardwood kraft pulp. Δ HexA denotes the amount of hexamonic acid groups removed in the acid treatment. Treatments were carried out at 10 % pulp consistency. The pH 1.5 corresponds to a charge of 2 % of sulfuric acid on pulp.

pH	Time (s)	Kappa number	Viscosity (ml/g)	Δ HexA (mmol/kg)
-	-	11.9	1005	-
1.5	1	10.7	975	9.7
1.5	2	9.5	919	14.9
1.5	4	8.4	834	30.9
3.5	1	11.2	985	4.6
3.5	2	10.5	961	11.8
3.5	4	8.7	955	29.2

EXHIBIT D

Table 3. Effect of acidic pretreatments on peroxide bleaching of a hardwood kraft pulp. Δ HexA denotes the amount of hexuronic acid groups removed in the acid treatment. Treatments were carried out at 10 % pulp consistency. The duration of treatment was 2 hours except for the treatment at pH 3.5 at 90 °C that lasted 6 hours.

A-stage pH	A-stage temperature (°C)	Kappa number before P-stage	Kappa number after P-stage	Δ HexA (mg/kg)
-	-	15.3	8.9	-
2	70	14.6	8.3	6.5
3.5	70	14.3	8.9	5.1
2	90	12.6	6.7	27.1
3.5	90	12.3	5.9	40.0

EXHIBIT E

Table 4. Effect of acidic pretreatment on peroxide bleaching of an oxygen bleached hardwood kraft pulp. ΔH_{oxA} denotes the amount of hexamethylenic acid groups removed in the acid treatment. Treatments were carried out at 10 % pulp consistency. The duration of treatment was 3 hours except for the treatment at pH 3.5 at 90 °C that lasted 6 hours.

A-stage pH	A-stage temperature (°C)	Kappa number before P-stage	Kappa number after P-stage	ΔH_{oxA} (mg/kg)
-	-	11.1	9.9	-
2	70	10.0	7.9	6.9
3.5	70	10.4	7.4	2.4
3	90	8.5	4.7	29.0
3.5	90	8.1	4.2	37.8